

hemoroidy.

- 1) Fumio Y. et al."A comparison of six models of commercially available automated immunologic fecal occult blood analyzers". The Journal of Clinical Laboratory Instruments and Reagents. 2006; 29(2): 121–30. [Japanese]
- 2) Carroll MRR. et al."Evaluation of quantitative faecal immunochemical tests for haemoglobin". Guildford Medical Device Evaluation Centre (GMEC), Guildford, UK, 2013; 73p.
- 3) Kazuo F. et al."3 kishu niyoru bensenketsujidoubunsekisouchi no hikakukentou". The Journal of Clinical Laboratory Instruments and Reagents. 2013; 36(5): 679–85. [Japanese]
- 4) Rapi S. et al."Effects of fecal sampling on preanalytical and analytical phases in quantitative fecal immunochemical tests for hemoglobin". Int J Biol Markers. 2017; 32(3): 261–6.
- 5) Randall E. et al."Evaluation of Hemo Tech NS-Plus system for use in a province-wide colorectal cancer screening program". Clinical Biochemistry. 2013; 46(4–5): 365–8.
- 6) Demian WLL. et al."Evaluation of the analytical performance of the novel NS-Prime system and examination of temperature stability of fecal transferrin compared with fecal hemoglobin as biomarkers in a colon cancer screening program". Practical Laboratory Medicine. 2015; 2: 29–36.
- 7) Røseth AG. et al."Assessment of the neutrophil dominating protein calprotectin in feces. A methodologic study". Scand J Gastroenterol. 1992; 27(9): 793–8.